

Expansions of Theta Functions and Applications

A. Raouf Chouikha *

Abstract

We prove that the classical theta function θ_4 may be expressed as

$$\theta_4(v, \tau) = \theta_4(0, \tau) \exp\left[-\sum_{p \geq 1} \sum_{k \geq 0} \frac{1}{p} \left(\frac{\sin \pi v}{(\sin(k + \frac{1}{2})\pi\tau)}\right)^{2p}\right].$$

We obtain an analogous expansion for the three other theta functions since they are related.

These results have several consequences. In particular, an expansion of the Weierstrass elliptic function will be derived. Actions of the modular group and other arithmetical properties will also be considered. Finally using a new expression for the Rogers-Ramanujan continued fraction we produce a simple proof of a Rogers identity.

Key words and phrases : theta functions, elliptic functions, q-series, Fourier series, continued fractions ¹

1 Introduction

The classical theta function is given by the doubly infinite sum

$$\theta(v, \tau) = \sum_{-\infty < n < +\infty} e^{i\pi n^2 \tau} e^{2i\pi n v}. \quad (1)$$

which locally converges uniformly for $v \in C$ the complex plane, and $\tau \in \mathcal{H}_+$ the upper-half plane of complex numbers with positive imaginary part.

A remarkable feature of the theta function is its dual nature. Indeed, when viewed as a function of v , we see it as an elliptic analog of the exponential function and it may be used in order to express elliptic functions. Since θ is periodic with period 1 and quasi-period τ . When considered as a function of τ , θ has modular properties with close connection to the partition function and the representation of integers as sums of squares.

The story of this function started with Euler, J. Bernoulli and Fourier who occasionally used the theta function among other closely related functions. However,

*Universite Paris 13 LAGA UMR 7539 Villetaneuse 93430, e-mail: chouikha@math.univ-paris13.fr

¹2000 *Mathematical Subject Classification* Primary: 33E05, 11F11; Secondary: 33E20, 34A20, 11F20

the systematic study of theta functions and their utilisation for the theory of elliptic functions is due to Jacobi. He remarked that theta can be represented by series whose convergence is very fast and which may be used for numerical computations of elliptic functions.

General properties of theta functions may be found in many references, among which [B], [M] or [W-W] contain a good description.

The above function θ is often called the θ_3 function of Jacobi. There are three other θ functions obtained by a change of characteristics

$$\begin{aligned}\theta_1(v, \tau) &= \sum_{-\infty \leq n \leq +\infty} (-1)^{\frac{n-1}{2}} q^{\left(\frac{n+1}{2}\right)^2} e^{i\pi(2n+1)v} = 2 \sum_{n \geq 0} (-1)^n q^{\left(\frac{n+1}{2}\right)^2} \sin((2n+1)\pi v) \\ \theta_2(v, \tau) &= \sum_{-\infty \leq n \leq +\infty} q^{\left(\frac{n+1}{2}\right)^2} e^{i\pi(2n+1)v} = 2 \sum_{n \geq 0} q^{\left(\frac{n+1}{2}\right)^2} \cos((2n+1)\pi v) \\ \theta_4(v, \tau) &= \sum_{-\infty \leq n \leq +\infty} (-1)^n q^{n^2} e^{i\pi 2nv} = 1 + 2 \sum_{n \geq 1} (-1)^n q^{n^2} \cos(2n\pi v)\end{aligned}$$

where $q = e^{i\pi\tau}$ verifies $|q| < 1$, and v is a complex number.

The four theta functions are related. Then we can choose any of these four theta functions and then define the remaining three in terms of the one chosen. All four theta functions are entire functions of v . All are periodic, the period of θ_1 and θ_2 is 2, and that of θ_3 and θ_4 is 1.

It is known that the zeros of θ_1 are $m + n\tau$, those of θ_2 are $\frac{1}{2} + m + n\tau$, those of θ_3 are $\frac{1}{2} + \tau + m + n\tau$ and those of θ_4 are $\frac{1}{2}\tau + m + n\tau$ where m, n are integers. From the knowledge of the zeros it is possible to obtain infinite products representing the theta functions, and from these products the partial fraction expansions of $\log \theta(v, \tau)$ follow.

The main goal of this paper is to state a trigonometric expansion of theta functions in powers of $\sin \pi v$ and to derive some applications. Namely

$$\log \frac{\theta_4(v, \tau)}{\theta_4(0, \tau)} = \sum_{p=1}^{\infty} c_{2p}(\tau) (\sin \pi v)^{2p}$$

where the coefficients c_{2p} , which depend on τ , have the following form

$$c_{2p}(\tau) = (-1)^{p+1} \frac{2^{2p+1}}{2p} \sum_{k \geq 0} \left(\frac{q^{2k+1}}{(1 - q^{2k+1})^2} \right)^p = -\frac{1}{p} \sum_{k \geq 0} \frac{1}{(\sin(k + \frac{1}{2})\pi\tau)^{2p}}$$

where $q = e^{i\pi\tau}$, $|q| < 1$. In the same way, we also obtain the expressions

$$\theta_3(v, \tau) = \theta_4(0, \tau) \exp \left[- \sum_{p \geq 1} \sum_{k \geq 0} \frac{1}{p} \left(\frac{\cos \pi v}{(\sin(k + \frac{1}{2})\pi\tau)} \right)^{2p} \right],$$

$$\theta_2(v, \tau) = \theta_4(0, \tau) \exp[i\pi(v + \frac{1}{4}\tau) - \sum_{p \geq 1} \sum_{k \geq 0} \frac{1}{p} \left(\frac{\cos \pi(v + \frac{1}{2}\tau)}{(\sin(k + \frac{1}{2})\pi\tau)} \right)^{2p}],$$

$$\theta_1(v, \tau) = \theta_4(0, \tau) \exp[i\pi(v - \frac{1}{2} + \frac{1}{4}\tau) - \sum_{p \geq 1} \sum_{k \geq 0} \frac{1}{p} \left(\frac{\sin \pi(v + \frac{1}{2}\tau)}{(\sin(k + \frac{1}{2})\pi\tau)} \right)^{2p}].$$

The above expressions for θ_4 and θ_3 are valid in the "strip" $|\frac{\sin \pi v}{\sin \frac{1}{2}\pi\tau}| < 1$, those relating to θ_2 and θ_1 are valid in the "strip" $|\frac{\sin \pi(v + \frac{1}{2}\tau)}{\sin \frac{1}{2}\pi\tau}| < 1$.

Some consequences of the above expansions may be deduced. In particular, the Weierstrass elliptic function $\wp(z)$ with primitive periods $2, 2\tau$ has an analogous expansion

$$\wp(z + \tau) = e_3 - \sum_{p \geq 1} [\sum_{k \geq 0} \frac{-2(2p+1)(\sin \frac{\pi z}{2})^{2p}}{(\sin(k + \frac{1}{2})\pi\tau)^{2p+2}} + \sum_{k \geq 0} \frac{4p(\sin \frac{\pi z}{2})^{2p}}{(\sin(k + \frac{1}{2})\pi\tau)^{2p}}].$$

We also examine properties of c_{2p} under the actions of the modular group and Landen or Gauss transformations. Some arithmetical applications using Lambert series will also be considered. An expression for the Rogers-Ramanujan continued fractions will be derived

$$R(q) = q^{2/5} e^{\sum_{p \geq 1, k \geq 0} \frac{q^{2p(k+1)}}{p} \left(\frac{(q^2 - q)^{2p} - (q^3 - 1)^{2p}}{(q^{5(2k+1)} - 1)^{2p}} \right)}.$$

The last one allows us to prove the well known Rogers identity

$$R(q) = q^{\frac{2}{5}} \prod_{k \geq 1} \frac{(1 - q^{10k-2})(1 - q^{10k-8})}{(1 - q^{10k-4})(1 - q^{10k-6})}.$$

2 Expansions of the theta functions

2.1 Classical expansions

Recall at first the following facts which will be useful in the sequel

Proposition 2-1 *The function $\theta_4(v, \tau)$ satisfies the triple product*

$$\theta_4(v, \tau) = \prod_{n \geq 1} (1 - q^{2n})(1 - q^{2n+1}e^{2i\pi v})(1 - q^{2n-1}e^{-2i\pi v})$$

where $q = e^{i\pi\tau}$, $v \in C$, $\tau \in \mathcal{H}_+$.

It is also known that $\log \theta_4$ has a Fourier series expansion, [B] or [W-W]

Proposition 2-2 *The theta function θ_4 has the following expansion*

$$\theta_4(v, \tau) = \theta_4(0, \tau) \exp\left[4 \sum_{n \geq 1} \frac{q^n}{1 - q^{2n}} \frac{(\sin n\pi v)^2}{n}\right].$$

Moreover, this expansion is valid for $\text{Im}\tau > 0$ and for $|\text{Im}v| < \frac{1}{2}\text{Im}\tau$.

We also obtain similar expansions for the other three theta functions, [B]. In [C] we proved the following

Proposition 2-3 *The theta function θ_4 may be expressed as*

$$\theta_4(v, \tau) = \theta_4(0, \tau) \exp\left[\sum_{p \geq 1} c_{2p}(\tau)(\sin \pi v)^{2p}\right]$$

where the coefficients c_{2p} satisfy the recursion relation

$$(2p+2)(2p+1)c_{2p+2}(\tau) - 4p^2c_{2p}(\tau) = \frac{4}{\pi^2}a_{2p}(\tau). \quad (2)$$

with the a_{2p} obeying the relation

$$\left(\frac{\pi}{2}\right)^2[(2p+2)(2p+1)a_{2p+2} - 4p^2a_{2p}] - 12e_3a_{2p} + 6 \sum_{0 < r < p} a_{2r}a_{2p-2r} = 0 \quad (3)$$

where $e_3 = -\frac{(\pi)^2}{12}[\theta_2^4(0, \tau) + \theta_3^4(0, \tau)]$ and $a_2 = \frac{\pi^2}{4}\theta_2^4(0, \tau)\theta_3^4(0, \tau)$. Moreover, this expansion is valid for $\text{Im}\tau > 0$ and for $|\sin \pi v| < 1$.

After eliminating the a_{2p} we get relations between c_{2p} coefficients only. The next result yields an analog version of the preceding

Proposition 2-4 *The theta function $\theta_4(v, \tau)$ may be expressed under the form*

$$\theta_4(v, \tau) = \theta_4(0, \tau) \exp\left[\sum_{p \geq 1} c_{2p}(\tau)(\sin \pi v)^{2p}\right]$$

where coefficients c_{2p} satisfy the system of recurrence equations for $p \geq 1$

$$(A) \quad \begin{cases} 4! \binom{2p+4}{4} c_{2p+4} = (2p+1)(2p+2)[(2p+2)(2p+3) + 4p^2 - c_0] c_{2p+2} \\ + (2p)^2 [c_0 - (2p)^2] c_{2p} - 6[(2p+1)(2p+2)c_{2p+2} - 2c_2 - \sum_{k=1}^p 2kc_{2k}]^2 \end{cases}$$

where $c_0 = -4[\theta_2^4(0, \tau) + \theta_3^4(0, \tau)]$, $c_2 = \frac{1}{2\pi^2} \frac{\theta_4''(0, \tau)}{\theta_4(0, \tau)}$ and

$$c_4 = \frac{1}{3}\theta_2^4(0, \tau)\theta_3^4(0, \tau) + \frac{1}{3}c_2.$$

Moreover, this expansion is valid for $\text{Im}\tau > 0$ and for $|\sin \pi v| < 1$.

Indeed, starting from (2) a calculation gives

$$\begin{cases} \frac{4}{\pi^2} [(2p+2)(2p+1)a_{2p+2} - 4p^2 a_{2p}] = (2p+1)(2p+2)(2p+3)(2p+4)c_{2p+4} - \\ (2p+1)(2p+2)[(2p+2)(2p+3) + 4p^2]c_{2p+2} + (2p)^4 c_{2p} \end{cases}.$$

We also have

$$\sum_{0 < r < p} a_{2r} a_{2p-2r} = \left(\frac{\pi}{2}\right)^4 [(2p+1)(2p+2)c_{2p+2} - c_2 - \sum_{k=1}^p c_{2k}]^2.$$

Thus, we obtain the following expressions connecting the c_{2p} coefficients

$$\begin{cases} (2p+1)(2p+2)(2p+3)(2p+4)c_{2p+4} - (2p+1)(2p+2)[(2p+2)(2p+3) \\ + 4p^2]c_{2p+2} + (2p)^4 c_{2p} + c_0[(2p+1)(2p+2)c_{2p+2} - \\ (2p)^2 c_{2p}] + 6[(2p+1)(2p+2)c_{2p+2} - 2c_2 - \sum_{k=1}^p 2kc_{2k}]^2 = 0 \end{cases}$$

which precisely are the recursion relations (A).

2.2 Main result

Now we are going to prove the following

Theorem 2-5 *Let $q = e^{i\pi\tau}$, $|q| < 1$. The coefficients c_{2p} defined by Proposition 2-4 may be expressed as*

$$c_{2p}(\tau) = -\frac{1}{p} \sum_{k \geq 0} \frac{1}{(\sin(k + \frac{1}{2})\pi\tau)^{2p}} = -\frac{1}{p} \sum_{k \geq 0} \left[\frac{(-4)q^{2k+1}}{(1 - q^{2k+1})^2} \right]^p.$$

Then, the theta functions have the following expansions

$$\theta_4(v, \tau) = \theta_4(0, \tau) \exp\left[-\sum_{p \geq 1} \sum_{k \geq 0} \frac{1}{p} \left(\frac{\sin \pi v}{(\sin(k + \frac{1}{2})\pi\tau)} \right)^{2p}\right],$$

$$\theta_3(v, \tau) = \theta_4(0, \tau) \exp\left[-\sum_{p \geq 1} \sum_{k \geq 0} \frac{1}{p} \left(\frac{\cos \pi v}{(\sin(k + \frac{1}{2})\pi\tau)} \right)^{2p}\right],$$

$$\theta_2(v, \tau) = \theta_4(0, \tau) \exp[i\pi(v + \frac{1}{4}\tau) - \sum_{p \geq 1} \sum_{k \geq 0} \frac{1}{p} \left(\frac{\cos \pi(v + \frac{1}{2}\tau)}{(\sin(k + \frac{1}{2})\pi\tau)} \right)^{2p}],$$

$$\theta_1(v, \tau) = \theta_4(0, \tau) \exp[i\pi(v - \frac{1}{2} + \frac{1}{4}\tau) - \sum_{p \geq 1} \sum_{k \geq 0} \frac{1}{p} \left(\frac{\sin \pi(v + \frac{1}{2}\tau)}{(\sin(k + \frac{1}{2})\pi\tau)} \right)^{2p}].$$

Moreover, the above expressions for θ_4 and θ_3 are valid in the "strip" $|\frac{\sin \pi v}{\sin \frac{1}{2}\pi\tau}| < 1$,

those for θ_2 and θ_1 are valid in the "strip" $|\frac{\sin \pi(v + \frac{1}{2}\tau)}{\sin \frac{1}{2}\pi\tau}| < 1$.

Proof of Theorem 2-5 At first, let us determine the coefficient $c_2(\tau)$.

Lemma 2-6 The coefficient $c_2(\tau)$ of the expansion of theta function given by Proposition 2-4 may be written under the forms

$$c_2(\tau) = -4 \frac{\sum_{n \geq 0} (-1)^n n^2 q^{n^2}}{1 + 2 \sum_{n \geq 1} (-1)^n q^{n^2}} = 4 \sum_{n \geq 1} \frac{q^{2n-1}}{(1 - q^{2n-1})^2} = 4 \sum_{n \geq 1} \frac{nq^n}{(1 - q^{2n})}$$

Indeed, from Propostion 2-4 one has

$$2\pi^2 c_2 = \frac{\theta_4''(0)}{\theta_4(0)}.$$

Furthermore, by the Fourier series expansion

$$\theta_4(v) = 1 + 2 \sum_{n \geq 0} (-1)^n e^{in^2\pi\tau} \cos(2n\pi v),$$

one deduces $\theta_4(0) = \sum_{n \geq 1} (-1)^n e^{in^2\pi\tau}$ and

$$\theta_4''(0) = -8\pi^2 \sum_{n \geq 1} (-1)^n n^2 e^{in^2\pi\tau}.$$

On the other hand, since we have (see [W-W] or [B] vol 3)

$$\frac{\theta_4''(0)}{\theta_4(0)} = 8\pi^2 \sum_{n \geq 1} \frac{q^{2n-1}}{(1 - q^{2n-1})^2}$$

we then get the second expression of c_2 , where $q = e^{i\pi\tau}$.

We shall compare different expansions for the theta functions. This will allow us to establish the expression of the c_{2p} coefficients.

By Proposition 2-2

$$\theta_4(v, \tau) = \theta_4(0, \tau) \exp[4 \sum_{n \geq 1} \frac{q^n}{1 - q^{2n}} \frac{(\sin n\pi v)^2}{n}]. \quad (4)$$

Comparing the latter with the expansion given by Proposition 2-4

$$\theta_4(v, \tau) = \theta_4(0, \tau) \exp\left[\sum_{p \geq 1} c_{2p}(\tau)(\sin \pi v)^{2p}\right]$$

one derives the following

Lemma 2-7 *The coefficients c_{2p} defined by Proposition 2-4 may be written as*

$$\begin{aligned} c_{2p}(\tau) &= (-1)^{p+1} 2^{2p} \left[\frac{q^p}{p(1-q^{2p})} + 2 \sum_{m > p} \binom{m+p-1}{m-p-1} \frac{q^m}{(m-p)(1-q^{2m})} \right] \\ &= (-1)^{p+1} \frac{2^{2p+1}}{(2p)!} \sum_{n \geq p} \frac{(n+p-1)!}{(n-p)!} \frac{q^n}{(1-q^{2n})}. \end{aligned}$$

In particular, we again find

$$c_2(\tau) = 4 \sum_{n \geq 1} \frac{nq^n}{(1-q^{2n})}.$$

Proof of Lemma 2-7 It is enough to express $\cos(2n\alpha)$ in terms of $\sin \alpha$ as

$$\begin{aligned} \cos(2n\alpha) &= \frac{(-1)^n}{2} (2 \sin \alpha)^{2n} + \sum_{k=0}^{n-1} (-1)^{n+k+1} \frac{2n}{k+1} \binom{2n-k+1}{k} (2 \sin \alpha)^{2n-2k-2} \\ &= n \sum_{0 \leq p \leq n} \frac{(-1)^p (n+p-1)!}{(2p)!(n-p)!} (2 \sin \alpha)^{2p} = \sum_{p=0}^n b_{2p,n} (\sin \alpha)^{2p}, \end{aligned}$$

where

$$b_{2n,n} = (-1)^n 2^{2n-1}; \quad \text{and} \quad b_{2p,n} = (-1)^p 2^{2p} \frac{n}{n-p} \binom{n+p-1}{n-p-1} \text{ for } n \neq p.$$

It can also be rewritten

$$b_{2p,n} = n \frac{(-1)^p (n+p-1)!}{(2p)!(n-p)!} 2^{2p} \quad \text{for } 0 \leq p \leq n.$$

Furthermore, we may use the fact

$$\log \frac{\theta_4(v, \tau)}{\theta_4(0, \tau)} = 2 \sum_{n \geq 1} \gamma_n - 2 \sum_{n \geq 1} \gamma_n \cos(2nv) = \sum_{p \geq 1} c_{2p}(\tau) (\sin \pi v)^{2p},$$

where $\gamma_n = \frac{q^n}{n(1-q^{2n})}$.

So, we obtain the relations

$$c_{2p} = -2 \sum_{n \geq p} \gamma_n b_{2p,n} = -(-1)^p 2^{2p} \left[\gamma_p b_{2p,p} + \sum_{n \geq p} \frac{q^n}{(n-p)(1-q^{2n})} \binom{n+p-1}{n-p-1} \right].$$

Therefore

$$c_{2p} = -2(-1)^p 2^{2p-1} \frac{q^p}{1-q^{2p}} + (-1)^{p+1} 2^{2p+1} \sum_{m>p} \binom{m+p-1}{m-p-1} \frac{q^m}{(m-p)(1-q^{2m})}.$$

i.e.,

$$c_{2p}(\tau) = (-1)^{p+1} \frac{2^{2p+1}}{(2p)!} \sum_{n \geq p} \frac{(n+p-1)!}{(n-p)!} \frac{q^n}{(1-q^{2n})}.$$

In particular, for $p = 1$,

$$c_2(\tau) = 2^2 \frac{q}{1-q^2} + 8 \sum_{m>1} \binom{m}{m-2} \frac{q^m}{(m-1)(1-q^{2m})},$$

and

$$c_2(\tau) = 2^2 \frac{q}{1-q^2} + 4 \sum_{m>1} \frac{mq^m}{(1-q^{2m})} = 4 \sum_{n \geq 1} \frac{nq^n}{(1-q^{2n})}.$$

These series can be expressed as

$$\begin{aligned} (-1)^{p+1} \frac{(2p)!}{2^{2p+1}} c_{2p}(\tau) &= \sum_{n \geq p} \frac{(n+p-1)!}{(n-p)!} \frac{q^n}{(1-q^{2n})} = \sum_{n \geq p} \frac{q^n (n+p-1)!}{(n-p)!} \sum_{k \geq 0} q^{2kn} \\ &= \sum_{k \geq 0, n \geq p} \frac{(n+p-1)!}{(n-p)!} q^{2kn+n} = \sum_{k \geq 0} q^{(2k+1)p} \sum_{n \geq p} \frac{(n+p-1)!}{(n-p)!} q^{(2k+1)(n-p)} \\ &= \sum_{k \geq 0} g_p(q^{2k+1}), \end{aligned}$$

where

$$g_p(z) = \sum_{n \geq p} \frac{(n+p-1)!}{(n-p)!} z^n = z^p \sum_{m \geq 0} \frac{(m+2p-1)!}{(m)!} q^m.$$

Now using the notation

$$(2p)_n = 2p(2p+1)\dots(2p+n-1), \quad \text{then} \quad (m+2p-1)! = (2p)_m (2p-1)!,$$

and one gets

$$\frac{1}{(2p-1)!} g_p(z) = z^p \sum_{m \geq 0} \frac{(2p)_m}{(m)!} q^m.$$

However, we know that

$$z^p \sum_{m \geq 0} \frac{(2p)_m}{(m)!} q^m = z^p {}_2F_1(2p, \alpha, \alpha, z) = \frac{z^p}{(1-z)^{2p}}.$$

$$g_p(\tau) = (2p-1)! \frac{z^p}{(1-z)^{2p}}.$$

As a result, we may derive the expression

$$c_{2p}(\tau) = (-1)^{p+1} \frac{2^{2p+1}}{(2p)!} \sum_{k \geq 0} g_p(q^{2k+1}) = (-1)^{p+1} \frac{2^{2p+1}}{2p} \sum_{k \geq 0} \frac{q^{(2k+1)p}}{(1 - q^{2k+1})^{2p}}.$$

$$(-1)^{p+1} \frac{(2p)!}{2^{2p+1}} c_{2p}(\tau) = (2p-1)! \sum_{k \geq 0} {}_2F_1(2p, \alpha, \alpha, q^{2k+1})$$

Finally,

$$c_{2p}(\tau) = (-1)^{p+1} \frac{2^{2p+1}}{2p} \sum_{k \geq 0} \left(\frac{q^{2k+1}}{(1 - q^{2k+1})^2} \right)^p = -\frac{1}{p} \sum_{k \geq 0} \frac{1}{(\sin(k + \frac{1}{2})\pi\tau)^{2p}}.$$

The proof of Theorem 2-5 has been achieved.

2.3 The link with elliptic functions

Now, consider the zeta function of Jacobi. It is defined by

$$Zn(z, k) = \frac{1}{2K} \frac{d}{dz} \log \theta_4(v, \tau),$$

where $v = \frac{z}{2K}$ and $K = 2 \int_0^{\frac{\pi}{2}} \frac{dx}{\sqrt{1-k^2 \sin^2 x}}$ is the complete elliptic integral of the first kind and the modulus is such that $0 < k < 1$.
We have

Corollary 2-8 *The zeta function of Jacobi has the following form*

$$Zn(z, k) = \frac{\pi}{2K} \sin(\pi 2v) \sum_{k \geq 0} \frac{1}{\sin^2(\pi v) - \sin^2(k + \frac{1}{2}\pi\tau)}$$

where $v = \frac{z}{2K}$ satisfies $|\sin \pi v| < |\sin(\frac{1}{2}\pi\tau)|$.

In particular, the logarithmic derivatives of theta functions can be written under the forms

$$\frac{\theta'_4(v, \tau)}{\theta_4(v, \tau)} = 4\pi \sin(\pi 2v) \sum_{k \geq 0} \frac{q^{2k+1}}{1 - 2q^{2k+1} \cos 2\pi v + q^{4k+2}}$$

$$\frac{\theta'_3(v, \tau)}{\theta_3(v, \tau)} = -4\pi \sin(\pi 2v) \sum_{k \geq 0} \frac{q^{2k+1}}{1 + 2q^{2k+1} \cos 2\pi v + q^{4k+2}}$$

$$\frac{\theta'_2(v, \tau)}{\theta_2(v, \tau)} = -\tan(\pi v) - 4\pi \sin(\pi 2v) \sum_{k \geq 0} \frac{q^{2k+2}}{1 + 2q^{2k+2} \cos 2\pi v + q^{4k+4}}$$

$$\frac{\theta'_1(v, \tau)}{\theta_1(v, \tau)} = \cot(\pi v) + 4\pi \sin(\pi 2v) \sum_{k \geq 0} \frac{q^{2k+2}}{1 - 2q^{2k+2} \cos 2\pi v + q^{4k+4}}.$$

Moreover, the equations for θ_1 and θ_2 are valid in the strip $|Imv| < Im\tau$, those for θ_3 and θ_4 are valid in the strip $|Imv| < \frac{1}{2}Im\tau$.

Indeed,

$$Zn(z, k) = \frac{1}{2K} \frac{d}{dz} \log \theta_4(v, \tau) = \frac{\pi}{2K} \sin(\pi 2v) \sum_{k \geq 0} \sum_{p \geq 1} \left(\frac{\sin \pi v}{(\sin(k + \frac{1}{2})\pi\tau)} \right)^{2p}.$$

Suppose the variable v satisfies

$$\left| \frac{\sin \pi v}{(\sin(\frac{1}{2})\pi\tau)} \right| < 1.$$

We then obtain

$$\begin{aligned} \left(\frac{\sin \pi v}{(\sin(k + \frac{1}{2})\pi\tau)} \right)^2 \sum_{p \geq 0} \left(\frac{\sin \pi v}{(\sin(k + \frac{1}{2})\pi\tau)} \right)^{2p} &= \frac{\left(\frac{\sin \pi v}{(\sin(k + \frac{1}{2})\pi\tau)} \right)^2}{1 - \left(\frac{\sin \pi v}{(\sin(k + \frac{1}{2})\pi\tau)} \right)^2} = \\ &= \frac{(\sin \pi v)^2}{(\sin(k + \frac{1}{2})\pi\tau)^2 - (\sin \pi v)^2}. \end{aligned}$$

Therefore, the result follows. The domain of convergence for these series may be extended to the strip $|Imv| < \frac{1}{2}Im\tau$ (see for example [W-W] page 489). Notice that the zeta function of Jacobi also has a Fourier expansion

$$Zn(z, k) = \frac{2\pi}{K} \sum_{n \geq 1} \frac{q^n}{1 - q^{2n}} \sin \frac{n\pi z}{K}.$$

Remark 2-10 (i) Concerning the convergence of the trigonometric series, notice that those pertaining to θ_3 and θ_4 could converge if $\cosh(Imv) < \sinh \frac{Im\tau}{2}$. Those applicable to θ_1 and θ_2 could converge if $|\coth(Imv)| < \tanh \frac{Im\tau}{2}$.

The first inequality implies $\left| \frac{\sin \pi v}{\sin \frac{1}{2}\pi\tau} \right| < 1$ and the second one implies $\left| \frac{\sin \pi(v + \frac{1}{2}\tau)}{\sin \frac{1}{2}\pi\tau} \right| < 1$.

(ii) Moreover, the above trigonometric expansions of theta functions which are very closed to Fourier series expansions seem to be new. We have not yet found an analog of these series (in this present form) nor an allusion to them in the classical literature concerning the theta functions. The only thing which may catch our attention is the connection of the above expansions of theta functions with the infinite products.

Indeed, The latter expressions given by Corollary 2-8 can also be deduced by logarithmic differentiation of the wellknown identities which yields the theta functions as infinite product. Therefore, the series expansions stated by Theorem

2-5 can also be deduced.
More precisely, from

$$\begin{aligned}\theta_1(v, \tau) &= 2q^{\frac{1}{4}} \sin(\pi z) \prod_{k \geq 0} (1 - q^{2k+2})(1 - 2q^{2k+2} \cos(2\pi v) + q^{4k+4}) \\ \theta_2(v, \tau) &= 2q^{\frac{1}{4}} \cos(\pi z) \prod_{k \geq 0} (1 - q^{2k+2})(1 + 2q^{2k+2} \cos(2\pi v) + q^{4k+4}) \\ \theta_3(v, \tau) &= \prod_{k \geq 0} (1 - q^{2k+2})(1 + 2q^{2k+1} \cos(2\pi v) + q^{4k+2}) \\ \theta_4(v, \tau) &= \prod_{k \geq 0} (1 - q^{2k+2})(1 - 2q^{2k+1} \cos(2\pi v) + q^{4k+2})\end{aligned}$$

we get the above expression of $\frac{\theta'_1(v, \tau)}{\theta_1(v, \tau)}, \frac{\theta'_2(v, \tau)}{\theta_2(v, \tau)}, \frac{\theta'_3(v, \tau)}{\theta_3(v, \tau)}, \frac{\theta'_4(v, \tau)}{\theta_4(v, \tau)}$, respectively (see [W-W] page 489).

However, we may notice that the previous infinite product representing the theta functions are valid in the entire v -plan.

Corollary 2-11 *Under the same hypotheses, the following expressions for ratios of theta functions hold*

$$\begin{aligned}\frac{\theta_1(v, \tau)}{\theta_2(v, \tau)} &= \exp\left[\sum_{p \geq 1} c_{2p}(\tau) [\sin^{2p} \pi(v + \frac{1}{2}\tau) - \cos^{2p} \pi(v + \frac{1}{2}\tau)]\right] \\ \frac{\theta_3(v, \tau)}{\theta_4(v, \tau)} &= \exp\left[\sum_{p \geq 1} c_{2p}(\tau) [\cos^{2p} \pi v - \sin^{2p} \pi v]\right].\end{aligned}$$

Corollary 2-12 *Under the same hypotheses, the following expression for the product of the theta functions holds*

$$\frac{\theta_2(v, \tau) \theta_3(v, \tau) \theta_4(v, \tau)}{\theta_4^3(0, \tau)} = e^{i\pi(v + \frac{1}{4}\tau)} \exp\left[\sum_{p \geq 1} c_{2p}(\tau) [\cos^{2p} \pi v + \sin^{2p} \pi v + \cos^{2p} \pi(v + \frac{\tau}{2})]\right].$$

In particular, we get

$$\theta'_1(0, \tau) = \pi \theta_2(0, \tau) \theta_3(0, \tau) \theta_4(0, \tau) = -\pi \theta_4^3(0, \tau) q^{\frac{1}{4}} \exp\left[\sum_{p \geq 1} c_{2p}(\tau) [1 + \cos^{2p} \pi \frac{\tau}{2}]\right].$$

The eta function of Dedekind may be expressed as

$$\eta(\tau) = 2^{-\frac{1}{3}} e^{i\pi\tau/12} \theta_4(0, \tau) \exp \sum_{p \geq 1} \frac{c_{2p}(\tau)}{3} [1 + \cos^{2p}(\pi \frac{\tau}{2})]$$

The proofs of the above results use the same techniques of [C].

3 Modular transformations

3.1 Some properties of the coefficients

The above coefficients $c_{2p}(\tau)$ are such that

$$c_{2p}(\tau + 2) = c_{2p}(\tau).$$

Furthermore, by the periodicity properties of the theta functions we may deduce some additional relations. More precisely, if the argument v is increased by τ , they are unaffected except for the multiplication by a simple factor. We get

$$\theta_1(v + \tau, \tau) = -(qe^{2i\pi v})^{-1}\theta_1(v, \tau)$$

$$\theta_2(v + \tau, \tau) = (qe^{2i\pi v})^{-1}\theta_2(v, \tau)$$

$$\theta_3(v + \tau, \tau) = (qe^{2i\pi v})^{-1}\theta_3(v, \tau)$$

$$\theta_4(v + \tau, \tau) = -(qe^{2i\pi v})^{-1}\theta_4(v, \tau)$$

We deduce the following

$$\sum_{p \geq 1} c_{2p}(\tau) [(\cos \pi v)^{2p} - (\sin \pi v)^{2p}] = i\pi + \sum_{p \geq 1} c_{2p}(\tau) [(\cos \pi(v + \tau))^{2p} - (\sin \pi(v + \tau))^{2p}].$$

$$\sum_{p \geq 1} c_{2p}(\tau) [(\cos \pi v)^{2p} - (\cos \pi(v + \tau))^{2p}] = i\pi(2v + \tau)$$

3.2 Actions of the modular group

Moreover, we have seen before (see [C]) that the double family of coefficients a_{2p} and c_{2p} , have analogous properties under the action of the modular group $\Gamma(1)$. For example, when we examine the theta relation $\theta_4(v + \frac{1}{2}, \tau + 1) = \theta_4(v, \tau)$, we find

$$c_{2p}(\tau + 1) = (-1)^p \sum_{k \geq p} \binom{k}{p} c_{2k}(\tau) \quad (5)$$

which is similar with the relation satisfied by the coefficients a_{2p} (see [C]). Equality (5), combined with System (A), permits to obtain other relations between the coefficients.

As consequences of Theorem 2-5, we have

Corollary 3-1 *Under the actions of the modular group $\tau \rightarrow \tau + 1$, and $\tau \rightarrow \frac{-1}{\tau}$, the coefficients become*

$$c_{2p}(\tau + 1) = -\frac{1}{p} \sum_{k \geq 0} \left[\frac{4q^{2k+1}}{(1 + q^{2k+1})^2} \right]^p.$$

$$\frac{2}{\tau^2} c_{2p}\left(\frac{-1}{\tau}\right) = -1 - (-1)^p \frac{2^{p+1}}{2p} \sum_{k \geq 1} \left[\frac{q^{2k}}{1 + q^{2k}} \right]^p.$$

Moreover, the following relation holds

$$\begin{aligned} (-1) \sum_{k \geq 1} k c_{2k}(\tau) &= \sum_{n \geq 1} \frac{n(-1)^n q^n}{(1 - q^{2n})}. \\ \sum_{p \geq 1} c_{2p}(\tau) \left[(\cos \pi \frac{\tau}{2})^{2p} \right] &= \sum_{p \geq 1} c_{2p}(\tau + 1) [1 + (\sin \pi \frac{\tau}{2})^{2p}]. \\ c_2(\tau + 1) &= -4 \sum_{k \geq 0} \frac{q^{2k+1}}{(1 + q^{2k+1})^2} = \frac{1}{2\pi^2} \frac{\theta_3''(0)}{\theta_3(0)}. \\ \frac{2}{\tau^2} c_2\left(\frac{-1}{\tau}\right) &= -1 - 8 \sum_{k \geq 1} \frac{q^{2k}}{1 + q^{2k}} = \frac{1}{2\pi^2} \frac{\theta_2''(0)}{\theta_2(0)}. \end{aligned}$$

3.3 Landen and Gauss transformations

Furthermore, using the Landen transformation we may obtain relations connecting $\theta(2v, 2\tau)$ and $\theta(v, \tau)$. More precisely, we get (see [B] or [W-W])

$$\theta_4(2v, 2\tau) = \frac{\theta_3(v, \tau) \theta_4(v, \tau)}{\theta_4(0, 2\tau)} = \theta_4(0, 2\tau) \exp\left[\sum_{p \geq 1} c_{2p}(2\tau) (\sin 2\pi v)^{2p}\right]. \quad (6)$$

From Proposition 2-4 and Theorem 2-5, one proves the following

Proposition 3-2 *The coefficients $c_{2p}(2\tau)$ satisfy the relation*

$$c_{2p}(2\tau + 1) = \sum_{k \geq 2p} 2^{-k} \binom{k}{2p} c_{2k}(\tau)$$

Indeed, we have seen that

$$\theta_3(v, \tau) = \theta_4(0, \tau) \exp\left[\sum_{p \geq 1} c_{2p}(\tau) (\cos \pi v)^{2p}\right].$$

Moreover, since $[\theta_4(0, 2\tau)]^2 = \theta_3(0, \tau) \theta_4(0, \tau)$ and $\log[\theta_3(0, \tau)] = \sum_p c_{2p}(\tau)$, then (6) implies

Lemma 3-3 *When v belongs to the strip $|\frac{\sin \pi v}{\sin \frac{1}{2} \pi \tau}| < 1$ the following relations hold*

$$\sum_{p \geq 1} c_{2p}(\tau) \left[(\sin \pi v)^{2p} + (\cos \pi v)^{2p} - 1 \right] = \sum_{p \geq 1} c_{2p}(2\tau) (\sin 2\pi v)^{2p}$$

$$\begin{aligned}
&= \sum_{p \geq 1} c_{2p}(2\tau + 1)(\cos 2\pi v)^{2p}. \\
\sum_{p \geq 1} c_{2p}(\tau) \left[(\sin \pi(v + \frac{1}{4}))^{2p} + (\cos \pi(v + \frac{1}{4}))^{2p} - 1 \right] &= \sum_{p \geq 1} c_{2p}(2\tau)(\cos 2\pi v)^{2p} \\
&= \sum_{p \geq 1} c_{2p}(2\tau + 1)(\sin 2\pi v)^{2p}.
\end{aligned}$$

In particular, we deduce the identities

$$\theta_3(2v, 2\tau) = \frac{\theta_3(v + \frac{1}{4}, \tau)\theta_3(v + \frac{1}{4}, \tau)}{\theta_4(0, 2\tau)} = \frac{\theta_4(v + \frac{1}{4}, \tau)\theta_4(v + \frac{1}{4}, \tau)}{\theta_4(0, 2\tau)}$$

Indeed, since $(\cos v)^{2n} = (\frac{1+\cos 2v}{2})^n = 2^{-n} \sum_{k=0}^{k=n} \binom{n}{k} (\cos 2v)^k$ then

$$\begin{aligned}
\sum_p 2^{-p} c_{2p}(\tau) \sum_{k=0}^{k=p} [(-1)^k \binom{p}{k} + \binom{p}{k}] (\cos 2v)^k - \sum c_{2p}(\tau) &= \\
&= \sum c_{2p}(2\tau + 1)(\cos 2v)^{2p},
\end{aligned}$$

thus

$$\sum_p 2^{-p} c_{2p}(\tau) \sum_{k=0}^{2k=p} \binom{p}{2k} (\cos 2v)^{2k} - \sum c_{2p}(\tau) = \sum c_{2p}(2\tau + 1)(\cos 2v)^{2p}.$$

As a result

$$\sum_k \left[\sum_{p \geq 2k} 2^{-p} \binom{p}{2k} c_{2p}(\tau) \right] (\cos 2v)^{2k} - \sum c_{2p}(\tau) = \sum c_{2p}(2\tau + 1)(\cos 2v)^{2p}.$$

Interchanging p and k , one gets

$$\sum_p \left[\sum_{k \geq 2p} 2^{-k} \binom{p}{2k} c_{2k}(\tau) \right] (\cos 2v)^{2p} - \sum c_{2p}(\tau) = \sum c_{2p}(2\tau + 1)(\cos 2v)^{2p}.$$

This proves Lemma 3-3 and Proposition 3-2.

Corollary 3-4 *The coefficients $c_{2p}(2\tau)$ also satisfy the equation*

$$c_{2p}(2\tau) = (-1)^p \sum_{m \geq 2p} \left[\sum_{k=p}^{2k=m} \binom{k}{p} \binom{m}{2k} \right] 2^{-m} c_{2m}(\tau)$$

Indeed, (5) implies $c_{2p}(\tau + 1) = (-1)^p \sum_{k \geq p} \binom{k}{p} c_{2k}(\tau)$ and Proposition 3-2 gives $c_{2p}(2\tau + 1) = \sum_{k \geq 2p} 2^{-k} \binom{k}{2p} c_{2k}(\tau)$.

Thus, combining these two equalities, one obtains

$$\begin{aligned} c_{2p}(2\tau) &= (-1)^p \sum_{k \geq p} \binom{k}{p} \left[\sum_{2k \geq m} \binom{m}{2k} \right] 2^{-m} c_{2m}(\tau) \\ &= (-1)^p \sum_{m \geq 2p} \left[\sum_{k=p}^{2k=m} \binom{k}{p} \binom{m}{2k} \right] 2^{-m} c_{2m}(\tau) \end{aligned}$$

Corollary 3-5 *The η function of Dedekind satisfies the following*

$$\frac{\eta(2\tau)}{\eta(\tau)} = 2^{-1/3} q^{\frac{1}{12}} \exp \sum_{p \geq 1} \frac{c_{2p}(\tau)}{3} \left[\cos^{2p} \frac{\pi\tau}{2} - 1 \right].$$

Indeed, this follows from the known identities

$$\frac{\eta^3(2\tau)}{\eta^3(\tau)} = \frac{\theta'_1(0, 2\tau)}{\theta'_1(0, \tau)} = \frac{1}{2} \frac{\theta_2(0, \tau)}{\sqrt{\theta_3(0, \tau)\theta_4(0, \tau)}}.$$

4 Some arithmetical properties

4.1 Transformations of higher order

One proved (Proposition (3-4) of [C1]) for $m \geq 1$ a modular equation satisfied by the coefficients

$$\frac{1}{2} \frac{q^m}{1 - q^{2m}} = m(-1)^{m+1} \sum_{p \geq m} 2^{-2p} \binom{2p}{p-m} c_{2p}(\tau) = \sum_{p \geq 1} 2^{-2p} \binom{2p}{p-1} c_{2p}(m\tau).$$

In particular, for $m = 2$ one again finds

$$-\sum_{p \geq 2} 2^{-2p} \binom{2p}{p-2} c_{2p}(\tau) = \sum_{p \geq 1} 2^{-2p} \binom{2p}{p-1} c_{2p}(2\tau)$$

In fact, for any positive integer k we get the identity

$$\begin{aligned} \frac{1}{2} \frac{q^m}{1 - q^{2m}} &= \sum_{p \geq m}^{k' \geq 0} \binom{2p}{p-1} \frac{(-1)^{p+1}}{p} \left(\frac{q^{2k'm+m}}{(1 - q^{2k'm+m})^2} \right)^p \\ &= k(-1)^{k+1} \sum_{p \geq k} 2^{-2p} \binom{2p}{p-k} c_{2p}\left(\frac{m}{k}\tau\right). \end{aligned}$$

Thus, we have

Proposition 4-1 *Let n and $k \geq 1$ be integers. The following identities hold*

$$\begin{aligned} k(-1)^{k+1} \sum_{p \geq k} 2^{-2p} \binom{2p}{p-k} c_{2p}(n\tau) &= n(-1)^{n+1} \sum_{p \geq n} 2^{-2p} \binom{2p}{p-n} c_{2p}(k\tau) = \\ \sum_{p \geq 1} 2^{-2p} \binom{2p}{p-1} c_{2p}(nk\tau) &= nk(-1)^{nk+1} \sum_{p \geq nk} 2^{-2p} \binom{2p}{p-nk} c_{2p}(\tau) \end{aligned}$$

4.2 Other expressions using Lambert series

Proposition 4-2 *The coefficients $c_{2p}(\tau)$ given by Theorem 2-5 may also be written under the form*

$$c_{2p}(\tau) = (-1)^{p+1} \frac{2^{2p+1}}{2p} \sum_{k \geq 0} \left(\frac{q^{2k+1}}{(1 - q^{2k+1})^2} \right)^p = (-1)^{p+1} \frac{2^{2p}}{2p!} \sum_{n \geq p} (A_n + B_n) q^n$$

where

$$A_n + B_n = \sum_{d|n, d \geq p} (1 + (-1)^{\frac{n}{d}-1}) \frac{(d+p-1)!}{(d-p)!}$$

Proof It is known that for $|z| < 1$ Lambert series can be written in different manners. In particular,

$$\sum_{n \geq 1} a_n \frac{z^n}{1 - z^n} = \sum_{n \geq 1} A_n z^n,$$

where

$$A_n = \sum_{d|n} a_d.$$

Moreover, using the Mobius function $\mu(n)$, we can express a_n in terms of A_n . Namely,

$$a_n = \sum_{d|n} \mu\left(\frac{n}{d}\right) A_d.$$

Furthermore, we may prove that

$$\sum_{n \geq 1} \frac{z^n}{1 + z^n} = \sum_{n \geq 1} B_n z^n,$$

where

$$B_n = \sum_{t|n} (-1)^{t-1} a_{\frac{n}{t}} = \sum_{d|n} (-1)^{\frac{n}{d}} a_d.$$

Let for any $p \geq 1$

$$c_{2p}(\tau) = (-1)^{p+1} \frac{2^{2p+1}}{(2p)!} \sum_{n \geq p} \frac{(n+p-1)!}{(n-p)!} \frac{q^n}{1 - q^{2n}}$$

Then, we may deduce

$$2(-1)^{p+1} \frac{2^{2p}}{(2p)!} \sum_{n \geq p} \frac{(n+p-1)!}{(n-p)!} \frac{z^n}{1 - z^{2n}} = \sum_{n \geq p} (A_n + B_n) z^n,$$

where

$$A_n + B_n = \sum_{d|n, d \geq p} (1 + (-1)^{\frac{n}{d}-1}) \frac{(d+p-1)!}{(d-p)!},$$

and $A_n = B_n = 0$ for $n < p$.

Remark 4-3 For $p = 1$, we get

$$c_2(\tau) = 4 \sum_{n \geq 1} \frac{nq^n}{(1 - q^{2n})}$$

but,

$$\frac{nq^n}{(1 - q^{2n})} = n(q^n + q^{3n} + q^{5n} + \dots)$$

The coefficient of q^m is n , if n divides m in such a way that $\frac{m}{n}$ is odd. Suppose $m = 2^k p$, p being odd. It implies that $n = 2^k d$, where d is a divisor of p .

Then, the coefficient of q^m in the expansion of $\frac{nq^n}{(1 - q^{2n})}$ is obviously $2^k \sum_{d|p} d$.

5 An expansion of $\wp(z)$

In this section we propose another interesting application of Theorem 2-5, one concerning a new type of expansion for elliptic functions. More precisely, we state for the elliptic Weierstrass function $\wp(z)$ an analogous expansion as we did for the θ function.

Recall that $\wp(z)$ which has primitive periods 2 and 2τ relative to g_2 and g_3 , may be written as

$$\wp(z) = \wp(z; 1, \tau) = \frac{1}{z^2} + \sum_{m,n} \left[\frac{1}{(z - 2m - 2n\tau)^2} - \frac{1}{(2m + 2n\tau)^2} \right]$$

The Weierstrass function $\wp(z)$ is related to the theta functions $\theta_i(v)$ where $v = \frac{z}{2\omega}$:

$$\wp(z) = \left(\frac{1}{2\omega}\right)^2 \left[-4\eta\omega - \frac{d^2 \log \theta_1(v)}{dv^2} \right]$$

with $\eta = -\frac{1}{12\omega} \frac{\theta_1'''(0)}{\theta_1'(0)}$. In the same way we have

$$\wp(z + \omega') = \left(\frac{1}{2\omega}\right)^2 \left[-4\eta\omega - \frac{d^2 \log \theta_4(v)}{dv^2} \right].$$

We proved (see [C]) that $\wp(z; 1, \tau)$ has the following expansion

$$\wp(z + \tau) = e_3 - \sum_{p \geq 1} a_{2p}(e_3) \left(\sin \frac{\pi z}{2} \right)^{2p}. \quad (7)$$

where $a_2(e_3) = \left(\frac{1}{\pi}\right)^2 (g_2 - 12e_3^2)$, $e_3 = \wp(\tau)$ and the other coefficients satisfy the recursion relations

$$\left(\frac{\pi}{2}\right)^2 [(2p+2)(2p+1)a_{2p+2} - 4p^2 a_{2p}] - 12e_3 a_{2p} + 6 \sum_{0 < r < p} a_{2r} a_{2p-2r} = 0. \quad (3)$$

Using relations (2) and (3) and Theorem 2-5 we shall give explicit expressions for the coefficients $a_{2p}(e_3) = a_{2p}(\tau)$

Theorem 5-1 *The Weierstrass elliptic function $\wp(z) = \wp(z; 1, \tau)$ with primitive periods 1 and τ may be expressed under the form*

$$\wp(z + \tau) = e_3 - \sum_{p \geq 1} a_{2p}(\tau) \left(\sin \frac{\pi z}{2} \right)^{2p}$$

where the coefficients are

$$a_{2p}(\tau) = -2(2p+1) \sum_{k \geq 0} \frac{1}{(\sin(k + \frac{1}{2})\pi\tau)^{2p+2}} + 4p \sum_{k \geq 0} \frac{1}{(\sin(k + \frac{1}{2})\pi\tau)^{2p}}.$$

Moreover, the above expression of $\wp(z)$ is valid in the strip $|\frac{\sin \pi v}{\sin \frac{1}{2}\pi\tau}| < 1$.

Indeed, the expression for $a_{2p}(\tau)$ is deduced from

$$(2p+2)(2p+1)c_{2p+2}(\tau) - 4p^2c_{2p}(\tau) = \frac{4\omega^2}{\pi^2}a_{2p}(\omega, \omega') = \frac{3}{\pi^2}a_{2p}(1, \tau) \quad (2)$$

and Theorem 4-1 since

$$c_{2p}(\tau) = -\frac{1}{p} \sum_{k \geq 0} \frac{1}{(\sin(k + \frac{1}{2})\pi\tau)^{2p}} = -\frac{1}{p} \sum_{k \geq 0} \left[\frac{(-4)q^{2k+1}}{(1 - q^{2k+1})^2} \right]^p.$$

Thus, $\wp(z)$ may be expressed as

$$\wp(z + \tau) = e_3 - \sum_{p \geq 1} \left[\sum_{k \geq 0} \frac{-2(2p+1)(\sin \frac{\pi z}{2})^{2p}}{(\sin(k + \frac{1}{2})\pi\tau)^{2p+2}} + \sum_{k \geq 0} \frac{4p(\sin \frac{\pi z}{2})^{2p}}{(\sin(k + \frac{1}{2})\pi\tau)^{2p}} \right].$$

Moreover, according to the addition theorem for $\wp(z)$, this function may be written as

$$\wp(z) = e_3 - \frac{g_2 - 12e_3^2}{2 \sum_{p \geq 1} \sum_{k \geq 0} \left[\frac{-2(2p+1)(\sin \frac{\pi z}{2})^{2p}}{(\sin(k + \frac{1}{2})\pi\tau)^{2p+2}} + \frac{4p(\sin \frac{\pi z}{2})^{2p}}{(\sin(k + \frac{1}{2})\pi\tau)^{2p}} \right]}.$$

6 The Rogers-Ramanujan continued fractions

For $|q| < 1$ the Rogers functions are defined by

$$G(q) = \sum_{n \geq 0} \frac{q^{2n^2}}{(q^2, q^2)_n}; \quad H(q) = \sum_{n \geq 0} \frac{q^{2n(n+1)}}{(q^2, q^2)_n}$$

where $q = e^{i\pi\tau}$; $(a, q)_n = \prod_{k=0}^{n-1} (1 - aq^k)$; $(a, q)_\infty = \lim_{n \rightarrow \infty} (a, q)_n$.
It is known that

$$G(q) = \frac{1}{(q^2, q^{10})_\infty (q^8, q^{10})_\infty}; \quad H(q) = \frac{1}{(q^4, q^{10})_\infty (q^6, q^{10})_\infty}.$$

The Rogers-Ramanujan continued fractions is defined by

$$R(q) = \frac{q^{2/5}}{1 + \frac{q^2}{1 + \frac{q^4}{1 + \frac{q^6}{1 + \dots}}}}.$$

It can be expressed in terms of q -series by

$$R(q) = q^{2/5} \frac{H(q)}{G(q)} = q^{2/5} \prod_{k \geq 1} \frac{(1 - q^{10k-2})(1 - q^{10k-8})}{(1 - q^{10k-4})(1 - q^{10k-6})}$$

We intend to give a simple proof of the the above Rogers identity.

Theorem 6-1 *The Rogers-Ramanujan continued fractions $R(q)$ for $q = e^{i\pi\tau}$, $|q| < 1$ may be expressed as*

$$\begin{aligned} R(q) &= \exp\left[i\pi\frac{2\tau}{5} - \sum_{p \geq 1} c_{2p}(5\tau) \left[\left(\sin \frac{\pi\tau}{2}\right)^{2p} - \left(\sin \frac{3\pi\tau}{2}\right)^{2p} \right]\right] \\ &= \exp\left[i\pi\frac{2\tau}{5} + \sum_{p \geq 1} \sum_{k \geq 0} \frac{1}{p} \frac{\left[\left(\sin \frac{\pi\tau}{2}\right)^{2p} - \left(\sin \frac{3\pi\tau}{2}\right)^{2p} \right]}{\left(\sin\left(k + \frac{1}{2}\right)5\pi\tau\right)^{2p}}\right]. \end{aligned}$$

Equivalently, in terms of q -series

$$R(q) = q^{2/5} e^{\sum_{p \geq 1, k \geq 0} \frac{q^{2p(k+1)}}{p} \left(\frac{(q^2 - q)^{2p} - (q^3 - 1)^{2p}}{(q^{5(2k+1)} - 1)^{2p}} \right)}.$$

Proof Consider the Ramanujan theta function

$$f(a, b) = \sum_{k \in \mathbb{Z}} a^{\frac{k(k+1)}{2}} b^{\frac{k(k-1)}{2}}$$

$$f(a, b) = (-a, ab)_\infty (-b, ab)_\infty (ab, ab)_\infty$$

with $|ab| < 1$. We denote here $(\alpha, \beta)_\infty = \prod_{i \geq 1} (1 - \alpha\beta^i)$. This function is related to the classical one

$$\theta_4(v, q) = f(-qe^{2i\pi v}, -qe^{-2i\pi v}).$$

Thus,

$$R(q) = q^{1/5} \frac{f(-q^2, -q^8)}{f(-q^4, -q^6)}.$$

One gets the relation

$$R(q) = q^{2/5} \frac{\theta_4(\frac{3}{2}\tau, 5\tau)}{\theta_4(\frac{1}{2}\tau, 5\tau)},$$

which implies the wanted expression of $R(q)$ since by Theorem 2-5

$$\theta_4(v, 5\tau) = \theta_4(0, 5\tau) \exp\left[-\sum_{p \geq 1} \sum_{k \geq 0} \frac{1}{p} \left(\frac{\sin \pi v}{\left(\sin\left(k + \frac{1}{2}\right)5\pi\tau\right)} \right)^{2p}\right].$$

Corollary 6-2 *$R(q)$ may also be expressed as*

$$R(q) = q^{2/5} \exp\left(\sum_{p \geq 1, k \geq 0} \frac{1}{p} \frac{\left[\left(\cos \frac{\pi\tau}{2}\right)^{2p} - \left(\cos \frac{3\pi\tau}{2}\right)^{2p} \right]}{\left(\cos\left(k + \frac{1}{2}\right)5\pi\tau\right)^{2p}}\right).$$

Equivalently,

$$R(q) = q^{2/5} e^{\sum_{p \geq 1, k \geq 0} \frac{q^{2p(k+1)}}{p} \left(\frac{(q^2 + q)^{2p} - (q^3 + 1)^{2p}}{(q^{5(2k+1)} + 1)^{2p}} \right)}.$$

Indeed, this is a direct consequence of the theta identity

$$\theta_4(v, \tau + 1) = \theta_3(v, \tau) = \theta_4(v + \frac{1}{2}, \tau).$$

Corollary 6-3 *Under the action of the modular group one has*

$$R(\tau + 1) = e^{2i\pi/5} R(\tau).$$

$$R(-1/\tau) = e^{i\pi/5 + \sum_{p \geq 1, k \geq 0} \frac{1}{p} \frac{[(\sin \frac{\pi(\tau-4)}{10})^{2p} - (\sin \frac{\pi(\tau-2)}{10})^{2p}]}{(\sin(k + \frac{1}{2})\pi\tau/5)^{2p}}}.$$

Indeed, the first identity is obtained from Corollary 6-2. Whereas, we may deduce the second one from

$$R(-1/\tau) = e^{i\pi/5} \frac{\theta_3(\frac{3+\tau}{10}, \tau/5)}{\theta_3(\frac{1+\tau}{10}, \tau/5)} = e^{i\pi/5} \frac{\theta_4(\frac{\tau-2}{10}, \tau/5)}{\theta_4(\frac{\tau-4}{10}, \tau/5)}.$$

Corollary 6-4 *Let $q = e^{i\pi\tau}$, $|q| < 1$. From the expression*

$$R(q) = q^{2/5} \exp \left(\sum_{p \geq 1, k \geq 0} \frac{1}{p} \frac{[(\sin \frac{\pi\tau}{2})^{2p} - (\sin \frac{3\pi\tau}{2})^{2p}]}{(\sin(k + \frac{1}{2})5\pi\tau)^{2p}} \right)$$

we may deduce the Rogers formula

$$R(q) = q^{\frac{2}{5}} \prod_{k \geq 1} \frac{(1 - q^{10k-2})(1 - q^{10k-8})}{(1 - q^{10k-4})(1 - q^{10k-6})}.$$

Proof Starting from the expansions

$$\begin{aligned} \sum_{p \geq 1} \frac{1}{p} \frac{(\sin \frac{\pi\tau}{2})^{2p}}{(\sin(k + \frac{1}{2})5\pi\tau)^{2p}} &= -\log(1 - \frac{(\sin \frac{\pi\tau}{2})^2}{(\sin(k + \frac{1}{2})5\pi\tau)^2}) \\ \sum_{p \geq 1} \frac{1}{p} \frac{(\sin \frac{3\pi\tau}{2})^{2p}}{(\sin(k + \frac{1}{2})5\pi\tau)^{2p}} &= -\log(1 - \frac{(\sin \frac{3\pi\tau}{2})^2}{(\sin(k + \frac{1}{2})5\pi\tau)^2}). \end{aligned}$$

We then derive the following expressions

$$\begin{aligned} R(q) &= q^{2/5} \prod_{k \geq 0} \frac{(\sin(k + \frac{1}{2})5\pi\tau)^2 - (\sin \frac{3\pi\tau}{2})^2}{(\sin(k + \frac{1}{2})5\pi\tau)^2 - (\sin \frac{\pi\tau}{2})^2} \\ R(q) &= q^{2/5} \prod_{k \geq 0} \frac{\cos(2k + 1)5\pi\tau - \cos 3\pi\tau}{\cos(2k + 1)5\pi\tau - \cos \pi\tau} \end{aligned}$$

$$R(q) = q^{2/5} \prod_{k \geq 0} \frac{[\sin(5k+4)\pi\tau][\sin(5k+1)\pi\tau]}{[\sin(5k+3)\pi\tau][\sin(5k+2)\pi\tau]}.$$

Thus,

$$R(q) = q^{2/5} \prod_{k \geq 0} \frac{(q^{10k+8} - 1)(q^{10k+2} - 1)q^{5k+3}q^{5k+2}}{q^{5k+4}q^{5k+1}(q^{10k+6} - 1)(q^{10k+4} - 1)}.$$

Remark 6-5 There are many other expressions of $R(q)$ in terms of theta functions. One of them given by Z.-G. Liu [L] may be interesting

$$R(q) = q^{-6/10} \frac{\theta_1(\tau, 5\tau)}{\theta_1(2\tau, 5\tau)}.$$

Indeed, since

$$\theta_1(v, \tau) = 2q^{1/4}(\sin \pi v)(q^2, q^2)_\infty (q^2 e^{-2i\pi v}, q^2)_\infty (q^2 e^{2i\pi v}, q^2)_\infty$$

then

$$\begin{aligned} \theta_1(\tau, 5\tau) &= iq^{1/4}(q^2, q^{10})_\infty (q^8, q^{10})_\infty (q^{10}, q^{10})_\infty; \\ \theta_1(2\tau, 5\tau) &= iq^{-3/4}(q^4, q^{10})_\infty (q^6, q^{10})_\infty (q^{10}, q^{10})_\infty. \end{aligned}$$

From which we deduce,

$$R(q) = q^{-6/10} \frac{(q^2, q^{10})_\infty (q^8, q^{10})_\infty}{(q^4, q^{10})_\infty (q^6, q^{10})_\infty} = q^{-6/10} \frac{\theta_1(\tau, 5\tau)}{\theta_1(2\tau, 5\tau)}.$$

The result follows from theorem 2-5 which asserts that

$$\theta_1(v, \tau) = \theta_4(0, \tau) \exp[i\pi(v - \frac{1}{2} + \frac{1}{4}\tau) - \sum_{p \geq 1} \sum_{k \geq 0} \frac{1}{p} \left(\frac{\sin \pi(v + \frac{1}{2}\tau)}{(\sin(k + \frac{1}{2})\pi\tau)} \right)^{2p}].$$

Thus,

$$R(q) = \exp[-i\pi \frac{8\tau}{5} + \sum_{p \geq 1} \sum_{k \geq 0} -\frac{1}{p} \frac{[(\sin \frac{7\pi\tau}{2})^{2p} - (\sin \frac{9\pi\tau}{2})^{2p}]}{(\sin(k + \frac{1}{2})5\pi\tau)^{2p}}].$$

Another equivalent expression may be obtained by the same techniques

$$R(q) = \exp[-i\pi \frac{8\tau}{5} + \sum_{p \geq 1} c_{2p}(5\tau + 1)[(\cos \frac{7\pi\tau}{2})^{2p} - (\cos \frac{9\pi\tau}{2})^{2p}].$$

References

- [A-L] P. Appell, E. Lacour, *Fonctions elliptiques et applications* Gauthiers-Villard ed., Paris (1922).
- [A] T.M. Apostol, *Modular functions and Dirichlet series in number theory* 2nd Ed. New-York; Springer-Verlag, (1997).
- [B] A.Erdélyi, W.Magnus, F.Oberhettinger, F.Tricomi, *Higher transcendental functions* Vol. I and III. Based on notes left by H. Bateman. Robert E. Krieger Publish. Co., Inc., Melbourne, Fla., (1981).
- [Be] R.E. Belmann, *A brief introduction to theta functions*, New York, Holt, Rinehart and Winston [1961]
- [C] R. Chouikha *On the expansions of elliptic functions and applications* in Algebraic methods and q-special functions, C.R.M. Proceedings and Lectures Notes, A.M.S., vol 22 , p.53-57, Providence, (1999).
- [C1] R. Chouikha *Sur des developements de fonctions elliptiques* Publ. Math., Fac. des Sc. de Besancon. Theorie des nombres, vol 8 , p. 1-9,(1989).
- [C2] R. Chouikha *Note on trigonometric expansions of theta functions* Proc. of the 6th OPSFA, (2002).
- [F] R. Fricke *Elliptische Funktionen.* Encycl. der Math. Wissensch., t.II, B.3, Teubner, Leipzig (1924).
- [L] Z.-G. Liu *Two theta functions identities and some Eisentein series identities of Ramanujan* Rocky Mountain J. of Math., vol 34, 2, p.713-732 (2004).
- [M] D. Mumford *Tata lectures on theta. vol II. Jacobian theta functions and differential equations* With C. Musili, M. Nori, E. Previato, M. Stillman and H. Umemura. Progress in Mathematics, 43. Birkhäuser, Boston, (1984).
- [W-W] E.T. Whittaker,G.N. Watson *A course of Modern Analysis* Cambridge (1963).